

Magnetic field correlations in a random flow with strong steady shear

I. Kolokolov, V. Lebedev, and G. Sizov
Landau Institute for Theoretical Physics RAS,
119334, Kosygina 2, Moscow, Russia,
and Moscow Institute of Physics and Technology.
 (Dated: October 29, 2010)

We analyze magnetic kinematic dynamo in a conducting fluid where the stationary shear flow is accompanied by relatively weak random velocity fluctuations. The diffusionless and diffusion regimes are described. The growth rates of the magnetic field moments are related to the statistical characteristics of the flow describing divergence of the Lagrangian trajectories. The magnetic field correlation functions are examined, we establish their growth rates and scaling behavior. General assertions are illustrated by explicit solution of the model where the velocity field is short-correlated in time.

PACS numbers: 47.35.Tv, 47.65.-d, 94.05.Lk, 95.30.Qd

I. INTRODUCTION

The subject of our work is the magnetic dynamo that is the magnetic field generation by hydrodynamic motions in a conducting medium. We theoretically investigate the effect in a conducting fluid (plasma, electrolyte) where a random hydrodynamic flow is excited. The principal example of such flow is hydrodynamic turbulence (see, e.g., Refs. [1, 2]) responsible for the magnetic field generation in different geophysical and astrophysical phenomena, see Refs. [3–10]. We consider the case where the magnetic field grows from small initial fluctuations and examine the evolution stage when the magnetic field is weak enough, so that one can neglect feedback from the magnetic field to the flow. The stage where the flow is independent of the magnetic field is called kinematic. The kinematic approach becomes invalid when the increasing magnetic field begins to affect the fluid motions essentially. In this case the velocity field is strongly influenced by the Lorentz force, so that the induction dynamics is no longer linear. In most cases that leads to saturation of the magnetic field fluctuations maintained by the hydrodynamic flow. Though the magnetic field cannot be described by a linear equation in the regime, the kinematic stage produces magnetic structures similar to those occurring at the saturation state, see, e.g., Ref. [11]. A possible explanation of the fact is related to strong intermittency of the magnetic field that implies that the feedback is concentrated in restricted space regions where the magnetic field is anomalously strong.

We assume that the random flow, exciting dynamo, is statistically homogeneous in space and time. Usually an additional assumption is made that the flow is statistically isotropic. If the velocity field is short-correlated in time then it is possible to derive closed equations for the magnetic induction correlation functions (see, e.g., Ref. [12]). The pair correlation function for the case has been analyzed in Refs. [13, 14]. The complete statistical description of the magnetic field for a short-correlated smooth statistically isotropic flow has been done in Ref. [15] where growth rates and structure of spatial correla-

tion functions were found. However, it is of interest to consider random flows with an average shear flow, that are widespread in astrophysical applications. Such flows are statistically anisotropic and need a special analysis. Here we examine the case where a steady shear flow is complemented by a relatively weak random component. We focus on the analysis of growth rates of moments of the magnetic field (magnetic induction), the degree of its anisotropy and on structure of the magnetic field correlation functions. We aim to relate the magnetic statistical characteristics to those of the flow, thus revealing the most universal features of the dynamo effect. The general assertions are illustrated by the model where the velocity field is short correlated in time, that can be solved analytically.

An additional motivation for our research comes from dynamics of polymer solutions that in many respects is similar to magneto-hydrodynamics, see Refs. [17, 18]. Particularly, we have in mind the so-called coil-stretch transition [19] (see also Refs. [20, 21]) that is an analog of the dynamo effect. A decade ago the elastic turbulence was discovered (see Refs. [22–24]) that is a chaotic hydrodynamic motion of polymer solutions, the elastic turbulence can be realized even at small Reynolds numbers as against the traditional hydrodynamic turbulence. The elastic turbulence is a natural frame for applying an extension of the dynamo theory to polymer solutions.

The behavior of the magnetic field moments at the kinematic stage in the presence of the strong shear flow was established in Ref. [25]. However, to examine the spacial structure of the magnetic field, one should know its correlation functions, that are studied in the present work in the framework of the same general scheme as in Ref. [25]. To verify our general predictions we examine the analytically solvable model where a short-correlated in time random flow is excited on background of the strong stationary shear flow.

The structure of our paper is as follows. In Section II we introduce basic relations needed to analyze the magnetic field correlations and dynamics. We present general dynamic equation, give its formally exact solution and

discuss statistical properties of quantities entering this solution. In Section III moments and correlation functions of the magnetic field are investigated. We relate its growth rates to the growth rates of the separation between two close fluid particles and establish principal space structure of the correlation functions. Section IV is devoted to the model where fluctuating component of the flow is short correlated in time. We establish the growth rates for the model and analyze in detail the pair correlation function. The obtained results are in agreement with our general assertions. In Section V we outline our main results and discuss their possible applications and extensions.

II. BASIC RELATIONS

We consider the magnetic field evolution in a conducting fluid (plasma or electrolyte) where hydrodynamic motions are excited. Then the magnetic field dynamics is governed by the following equation (see, e.g., Ref. [26])

$$\partial_t \mathbf{B} = (\mathbf{B} \cdot \nabla) \mathbf{v} - (\mathbf{v} \cdot \nabla) \mathbf{B} + \kappa \nabla^2 \mathbf{B}, \quad (2.1)$$

where $\mathbf{B}$ is the magnetic induction, $\mathbf{v}$ is the flow velocity and κ is the magneto-diffusion coefficient, inversely proportional to the electrical conductivity of the medium. The flow is assumed to be incompressible, $\nabla \cdot \mathbf{v} = 0$. We also assume that the magneto-diffusion term in Eq. (2.1) is small in comparison with those related to the flow. We consider the case where the magnetic field is relatively weak and, therefore, its feedback to the flow is negligible. Then the relation (2.1) is a linear equation determining the magnetic field evolution in a prescribed velocity field, this regime is called kinematic.

The hydrodynamic motion excited in the fluid is assumed to be random (turbulent) and the velocity statistics is assumed to be homogeneous in space and time. We examine the magnetic field growth from initial weak fluctuations distributed statistically homogeneously in space at the initial time $t = 0$. The correlation length of the initial fluctuations l is assumed to be smaller than the velocity correlation length η . If we consider hydrodynamic turbulence then a role of the velocity correlation length is played by the Kolmogorov scale. At scales less than η the velocity field $\mathbf{v}$ can be treated as smooth. The magnetic growth (dynamo) can be characterized by moments of the magnetic induction that exponentially increase over time t :

$$\langle |\mathbf{B}(t)|^{2n} \rangle \propto \exp(\gamma_n t), \quad (2.2)$$

Here angular brackets mean averaging over space. The exponential laws (2.2) are characteristic of the kinematic dynamo since the equation (2.1) is linear in the magnetic induction $\mathbf{B}$ in this case.

One of our goal is to express the growth rates (increments) γ_n in Eq. (2.2) via statistical characteristics of the flow. The natural measure for the growth rates γ_n is the

so-called Lyapunov exponent of the flow, λ , that is equal to the average logarithmic divergence rate of close fluid particles. A special question concerns an n -dependence of γ_n . If the magnetic induction statistics is Gaussian then $\gamma_n \propto n$. Deviations from the linear law signal about intermittency of the magnetic field. The intermittency implies that high moments of the magnetic field are determined by rare strong fluctuations.

There are two different regimes of the kinematic magnetic field growth. The first regime is realized if all characteristic scales of the magnetic field are much larger than the magnetic diffusion length r_d , $r_d = \sqrt{\kappa/\lambda}$. The assumed smallness of the diffusion coefficient implies the inequality $\eta \gg r_d$. We assume also $l \gg r_d$, then the diffusion term in Eq. (2.1) is negligible at the first stage of the magnetic evolution that we call diffusionless. The magnetic force lines are deformed by the flow without reconnections in this regime. However, distortions of the magnetic field by the flow inevitably lead to producing scales of order r_d in the field. After that the magnetic diffusion is switched on that admits reconnections. This second (diffusion) stage is characterized by the growth rates different from ones describing the diffusionless regime.

Let us present a qualitative picture explaining the magnetic field evolution at the kinematic stage. The initial magnetic field distribution in space can be thought as an ensemble of blobs of sizes $\sim l$. Then the blobs are distorted by the flow being stretched in one direction and compressed in another direction. In the isotropic case directions of the stretching and compression vary chaotically in space and time whereas in our case they are attached to the shear flow: the blobs are stretched mainly along the shear velocity and are compressed in the direction of the shear velocity gradient. At the first (diffusionless) stage the blobs are deformed without intersections and the magnetic field induction grows like a separation between close fluid particles since the equation (2.1) at $\kappa = 0$ coincides with the equation for the separations.

The diffusionless stage finishes when the characteristic blob width diminishes down to the diffusion length r_d . Then the diffusion is switched on that leads to two effects. First, the diffusion prevents further shrinking the blob widths, so that they remain of the order of r_d , whereas the blobs continue to be stretched in the direction of the shear velocity. Second, due to reconnections of the magnetic force lines admitted by diffusion the blobs start to overlap. As a result, new blobs of a characteristic longitudinal size η are formed, see Fig.1. The magnetic induction in such blobs can be found by averaging the induction of a large number N of initial blobs, the number N grows exponentially as time runs. Averaging over the large number of random variables leads to appearing an exponentially small factor $\sim 1/\sqrt{N}$ in the amplitude of the magnetic induction. Besides, the amplitudes of the initial blobs remain to increase over time as a separation between fluid particles. We conclude that at this second (diffusive) stage the magnetic field is still growing




FIG. 1: Sketch of typical magnetic blobs during the diffusive kinematic stage.

exponentially over time but slower than at the first stage.

We consider the case where the steady shear constituent of the flow is much stronger than the random one. Quantitatively, the condition is written as the inequality $s \gg \lambda$ where s is the shear rate. Indeed, the Lyapunov exponent in a pure shear flow is zero, and its non-zero value is associated with presence of the relatively weak random constituent of the flow. The distorted magnetic blobs are elongated mainly along the shear velocity. However, they are tilted with respect to the velocity direction due to presence of the random velocity component, see Fig. 1. The tilt possesses the same dynamics as the direction of the polymer stretching in the same flow, see Ref. [21]. Therefore the tilt angle ϕ , see Fig. 1, can be estimated as $\phi \sim \lambda/s$. The tilt angle determines typical ratio of the magnetic field components $B_y/B_x \sim \lambda/s \ll 1$ where the X -axis is directed along the shear velocity that varies along the Y -axis. Thus, the ratio s/λ characterizes an anisotropy degree of the magnetic field.

A. Lagrangian dynamics

To analyze moments and correlation functions of the magnetic induction we need a solution of the magnetodynamic equation (2.1) for the induction field $\mathbf{B}(t)$ in terms of its initial value $\mathcal{B}$, $\mathcal{B} = \mathbf{B}(0)$. We use here a generalization of the scheme proposed in Ref. [27] that exploits the Lagrangian approach to the fluid motion. Passing to the Lagrangian frame, that moves with the fluid particles, one excludes the advection term in Eq. (2.1). Next, instead of solving the equation with the Laplace operator ∇^2 it is convenient to introduce Langevin forces ξ mimicing the magnetic diffusion. Then we pass to the following stochastic equation

$$\partial_t \mathbf{R} = \mathbf{v}(t, \mathbf{R}) + \xi, \quad (2.3)$$

describing Lagrangian trajectories disturbed by the Langevin forces. The latter can be treated as white noise

characterized by its pair correlation function

$$\langle \xi_i(t_1) \xi_j(t_2) \rangle = 2\kappa \delta_{ij} \delta(t_1 - t_2), \quad (2.4)$$

where κ is the diffusion coefficient figuring in Eq. (2.1).

Rewriting the equation (2.1) in terms of the quantity $\mathbf{R}(t)$ one finds then the following formally exact solution of Eq. (2.1)

$$\mathbf{B}(t, \mathbf{r}) = \left[\hat{W}(t) \mathcal{B}[\mathbf{R}(0)] \right], \quad (2.5)$$

where floors mean averaging over the ξ -statistics, in accordance with Eq. (2.4). To find the quantity $\mathbf{R}(0)$ one should solve the equation (2.3) on the time interval $(0, t)$ with the boundary condition $\mathbf{R}(t) = \mathbf{r}$, posed at the final time. By other words, one should track the magnetic field back in time along the disturbed Lagrangian trajectories. The matrix $\hat{W}(t)$ in Eq. (2.5) is a chronologically ordered exponent

$$\hat{W}(t) = \text{T exp} \left\{ \int_0^t dt' \hat{\Sigma}(t') \right\}, \quad (2.6)$$

where $\hat{\Sigma}(t)$ is the velocity gradients matrix, $\Sigma_{ji} = \partial_i v_j$, taken at the time t and in the spacial point $\mathbf{R}(t)$. The matrix $\hat{W}$, that we call an evolution matrix, can be treated as a solution of the equation $\partial_t \hat{W} = \hat{\Sigma} \hat{W}$ with the initial condition $\hat{W}(0) = 1$.

The evolution matrix $\hat{W}$ has some general properties that follow from the definition (2.6). The determinant of the matrix $\hat{W}$ is equal to unity, since the velocity gradient matrix $\hat{\Sigma}$ is traceless, $\text{tr} \hat{\Sigma} = 0$, that, in turn, is a consequence of the incompressibility condition $\nabla \cdot \mathbf{v} = 0$. Therefore $W_1 W_2 W_3 = 1$, where W_1, W_2, W_3 are eigen values of the matrix $\hat{W}$. All the eigen values are positive since they are positive initially ($W_1 = W_2 = W_3 = 1$ at $t = 0$) and cannot turn to zero because of the same relation $W_1 W_2 W_3 = 1$. Let us order the eigen values in accordance with $W_1 > W_2 > W_3$, then $W_1 > 1$ and $W_3 < 1$. At times $t \gg \lambda^{-1}$, we are interested in, typical values of $\ln W_1, \ln W_3$ can be estimated as $\pm \lambda t$, therefore W_1 is exponentially large whereas W_3 is exponentially small. An estimation for W_2 depends on details of the flow statistics. Any case, $W_1 \gg W_2 \gg W_3$ at times $t \gg \lambda^{-1}$.

In the framework of the proposed formalism, correlation functions of the magnetic field $\mathbf{B}$ have to be calculated by averaging products of the factors (2.5) taken at the respective points over the statistics of the noises ξ , besides averaging over space. Thus, say, the simultaneous correlation function,

$$F_{2n, i \dots j}(\mathbf{r}_1, \dots, \mathbf{r}_{2n}) = \langle B_i(\mathbf{r}_1) \dots B_j(\mathbf{r}_{2n}) \rangle, \quad (2.7)$$

has to be calculated in two steps. First, one should substitute the expression (2.5) into the right-hand side of Eq. (2.7) and then average the resulting product over the ξ -statistics given by Eq. (2.4), this averaging catches diffusion effects. Let us underline that the fields ξ have to

be treated as independent for all $2n$ factors in the product. Second, one should average the result over space. An averaging over scales $\lesssim \eta$ (traced back to the initial time) gives statistics of the initial magnetic field fluctuations, and averaging over scales $\gtrsim \eta$ counts different realizations of $\hat{\Sigma}$. Therefore the latter is equivalent to averaging over the velocity statistics. This logic was realized for the isotropic random flow in Ref. [15].

In the diffusionless regime, realized at $t \ll \lambda^{-1} \ln(l/r_d)$, one can neglect diffusion effects. Then at calculating the moment $\langle |\mathbf{B}|^{2n} \rangle$ one can take a product of the identical factors (2.5) where $\mathbf{R}$ is simply a Lagrangian trajectory terminated at the point $\mathbf{r}$ at time t . Then $|\mathbf{B}(\mathbf{r})|^{2n} \approx W_1^{2n} |\mathcal{B}|^{2n}$ where $\mathcal{B}$ is taken at the origin of the Lagrangian trajectory. Here just the factor W_1^{2n} is responsible for the exponential growth of the moments, and therefore, we can restrict ourselves to the estimation $[|\mathbf{B}(\mathbf{r})|^{2n}] \sim W_1^{2n} \mathcal{B}_0^{2n}$ where $\mathcal{B}_0$ is the characteristic value of the initial magnetic field fluctuations. In the diffusion regime, realized at $t \gg \lambda^{-1} \ln(l/r_d)$, the situation is a bit more complicated.

Let us first consider the second moment. Then we deal with two trajectories, $\mathbf{R}$ and $\mathbf{R}'$, terminating at the same point $\mathbf{r}$ at time t , but characterized by independent noises ξ and ξ' . The second moment can be written as the following average

$$\langle (\mathbf{B})^2 \rangle = \left\langle \left[\mathcal{B}^T[\mathbf{R}(0)] \hat{W}^T \hat{W}' \mathcal{B}[\mathbf{R}'(0)] \right] \right\rangle, \quad (2.8)$$

where the subscript T means transposition of the object. An appreciable contribution to the second moment is related to the trajectories with $|\mathbf{R}(0) - \mathbf{R}'(0)| \lesssim l$. Since $|\mathbf{R}(0) - \mathbf{R}'(0)| \ll \eta$ and $|\mathbf{R}(t) - \mathbf{R}'(t)| = 0$, the difference $\Delta \mathbf{R} = \mathbf{R} - \mathbf{R}'$ stays to be much less than η at any time from the interval $(0, t)$ for such event. Then we obtain from Eq. (2.3) expanding the velocity up to linear in $\Delta \mathbf{R}$ terms

$$\partial_t \Delta \mathbf{R} = \hat{\Sigma} \Delta \mathbf{R} + \xi - \xi', \quad (2.9)$$

where $\hat{\Sigma}$ can be taken at the point $\mathbf{R}$ or $\mathbf{R}'$, no difference. A solution of Eq. (2.9), equal to zero at time $t' = t$, is written as

$$\Delta \mathbf{R}(t') = -\hat{W}(t') \int_{t'}^t dt_1 \hat{W}^{-1}(t_1) [\xi(t_1) - \xi'(t_1)]. \quad (2.10)$$

To calculate the second moment we should know the $\Delta \mathbf{R}(0)$ -statistics. Since the separation $\Delta \mathbf{R}(0)$ is a linear combination of ξ, ξ' it should be treated as a Gaussian variable at averaging over the ξ -statistics, then its probability distribution function is completely determined by the matrix $\hat{M}$:

$$M_{ij} = [\Delta R_i(0) \Delta R_j(0)] = 4\kappa \int_0^t dt_1 W_{ik}^{-1} W_{jk}^{-1}, \quad (2.11)$$

the expression for $\hat{M}$ is derived from Eqs. (2.4, 2.10). The matrix $\hat{M}$ is symmetric, and its eigen values are positive.

Let us designate the eigen values as m_1^2, m_2^2, m_3^2 and order the values as $m_1 > m_2 > m_3$, the inequalities become strong, $m_1 \gg m_2 \gg m_3$, if $\lambda t \gg 1$. Let us stress that the directions of the eigen vectors of the matrix $\hat{M}$ are “frozen” at $\lambda t \gg 1$ [29–31]. Then the integral determining m_1 is gained at $t - t_1 \sim \lambda^{-1}$ and we arrive at the estimation $m_1 \sim r_d W_3^{-1}$. The integral determining m_3 is gained at $t_1 \sim \lambda^{-1}$ and therefore $m_3 \sim r_d$. An estimation for m_2 depends on the time dependence of W_2 . If W_2 increases then m_2 remains of the order of r_d whereas it grows like $m_2 \sim r_d W_2^{-1}$ if W_2 decreases.

Now we find a probability that $\Delta \mathbf{R}(0)$ is less than l in the diffusion regime, when $t \gg \lambda^{-1} \ln(l/r_d)$. One can think in terms of the components of $\Delta \mathbf{R}(0)$ in the basis attached to the eigen vectors of the matrix $\hat{M}$. Since $m_1 \gg l$ then a probability that the first component of $\Delta \mathbf{R}(0)$ is less than l is estimated as $l/m_1 \sim (l/r_d) W_3$. If W_2 grows as time runs then both, m_2 and m_3 , are of the order of r_d and therefore a probability that the second and the third components of $\Delta \mathbf{R}(0)$ are less than l , is close to unity. Then we find from Eq. (2.8)

$$[|\mathbf{B}|^2] \sim \mathcal{B}_0^2 (l/r_d) W_1 W_2^{-1}, \quad (2.12)$$

where $\mathcal{B}_0$ is a characteristic value of the initial magnetic field fluctuations and we used the relation $W_1 W_2 W_3 = 1$.

The situation with the decreasing W_2 is slightly different. In this case $m_2 \gg l$ at the diffusive stage and there appears an additional small probability that the second component $\Delta \mathbf{R}(0)$ is less than l , the probability can be estimated as $l/m_2 \sim (l/r_d) W_2$. Then one would obtain $[|\mathbf{B}|^2] \sim \mathcal{B}_0^2 W_1 (l/r_d)^2$, instead of Eq. (2.12). However, an integration over space (at the next step of averaging) kills the leading term due to the solenoidal nature of the magnetic field $\mathbf{B}$. Therefore one has to take into account the next term in the probability distribution of $\Delta \mathbf{R}_2(0)$ that gives an extra small factor $(l/m_2)^2$ in the probability. Thus, we arrive at

$$[|\mathbf{B}|^2] \sim \mathcal{B}_0^2 (l/r_d)^4 W_1 W_2^2. \quad (2.13)$$

Note that the expressions (2.12, 2.13) are equivalent to ones obtained in the Fourier representation for the statistically isotropic case in Ref. [15]. However, the expressions (2.12, 2.13), written for real space, are correct for the anisotropic problem (we are investigating) as well, and are in fact more suitable for the problem.

Let us turn to the high moments. One can recognize that a principal contribution to the average $[|\mathbf{B}|^{2n}]$ is produced by configurations where the $2n$ points $\mathbf{R}_\alpha(0)$ are divided into n pairs with separations $\lesssim l$ in each pair. Because of the independence of the white noises ξ_α , the probability of such event can be estimated as a product of probabilities for the second moment, that is

$$[|\mathbf{B}|^{2n}] \sim [|\mathbf{B}|^2]^n, \quad (2.14)$$

where the second moment is given by Eq. (2.12) or Eq. (2.13). We have ignored a combinatoric factor in Eq. (2.14) being interested in the time dependence of the moments.

B. Evolution matrix

The next step in finding the magnetic field moments is averaging over the velocity statistics. Before doing it we should establish statistical properties of the evolution matrix (2.6). Some universal properties of such matrices, that can be treated as products of a large number of random matrices, are well established, see Refs. [29–31], the properties are revealed at $t \gg \lambda^{-1}$. However, we examine a strongly anisotropic case, with steady shear flow dominating. That forced us to modify the consideration of Ref. [15] where the isotropic case (statistically isotropic flow) was investigated.

For the anisotropic problem, it is convenient to use the Gaussian decomposition of the evolution matrix $\hat{W} = \hat{T}_L \hat{\Delta} \hat{T}_R$, where $\hat{T}_L$ and $\hat{T}_R$ are triangle matrices,

$$\hat{T}_L = \begin{pmatrix} 1 & \chi & \chi_1 \\ 0 & 1 & \chi_2 \\ 0 & 0 & 1 \end{pmatrix}, \quad \hat{T}_R = \begin{pmatrix} 1 & 0 & 0 \\ \zeta_1 & 1 & 0 \\ \zeta_2 & \zeta_3 & 1 \end{pmatrix}, \quad (2.15)$$

and $\hat{\Delta}$ is a diagonal matrix. Since both triangle matrices, $\hat{T}_L$ and $\hat{T}_R$, have unit determinants, the determinant of $\hat{\Delta}$ is equal to unity as well.

The matrices are written in the reference frame attached to the shear flow: the axis X is directed along the shear velocity and the axis Y is directed along the shear velocity gradient. Therefore the shear velocity is written as $v_x = sy$ where s is the shear rate. For our flow, which is composed of the steady shear flow and a random component, the matrix of the velocity gradients $\Sigma_{ji} = \partial_i v_j$ is a sum of two terms related to the shear and the random components of the flow:

$$\Sigma_{ji}(t) = s\delta_{jx}\delta_{iy} + \sigma_{ji}(t). \quad (2.16)$$

The random matrix σ_{ji} is zero in average and should be characterized in terms of its correlation functions. The trace of the matrix is zero, $\text{tr} \hat{\sigma} = 0$ (due to the flow incompressibility). Let us remind that the Lyapunov exponent λ of a purely shear flow is equal to zero. Therefore λ is sensitive to $\hat{\sigma}$ though the random flow is weaker than the steady one.

Substituting the decomposition $\hat{W} = \hat{T}_L \hat{\Delta} \hat{T}_R$ into the evolution equation $\partial_t \hat{W} = \hat{\Sigma} \hat{W}$, one finds

$$\hat{T}_L^{-1} \hat{\Sigma} \hat{T}_L = \hat{T}_L^{-1} \partial_t \hat{T}_L + \partial_t \hat{\Delta} \hat{\Delta}^{-1} + \hat{\Delta} \partial_t \hat{T}_R \hat{T}_R^{-1} \hat{\Delta}^{-1}. \quad (2.17)$$

The terms in the right-hand side of Eq. (2.17) are the left-off-diagonal matrix, the diagonal matrix, and the right-off-diagonal matrix, accordingly. Therefore, one obtains a closed (non-linear) equation for the matrix $\hat{T}_L$ that leads to homogeneous in time statistics of the matrix. Next, we obtain for components of the diagonal matrix $\partial_t \hat{\Delta} \hat{\Delta}^{-1}$ expressions that are random variables with statistics homogeneous in time. Therefore $\ln \Delta_1$, $\ln \Delta_2$, and $\ln \Delta_3$ (where Δ_i are eigen values of the matrix $\hat{\Delta}$) are subjects of the central limit theorem. Typically, the variables behave linear in time t with coefficients of the

order of λ . The situation with the matrix $\hat{T}_R$ is slightly more complicated since there are the exponential factors in the last term of Eq. (2.17). Therefore some components of the matrix $\hat{T}_R$ behave exponentially in time like the factors.

Based on the leading role of the shear term in the expression (2.16), one obtains from Eq. (2.17) for $\hat{T}_L$ a hierarchy $\chi \gg \chi_1 \gg \chi_2$. Therefore in the principal approximation in λ/s the only component, σ_{yx} , is relevant and the equation for the matrix $\hat{T}_L$ is reduced to a single equation for the component χ :

$$\partial_t \chi = s - \chi^2 \sigma, \quad (2.18)$$

where $\sigma \equiv \sigma_{yx}$. We conclude that the variable χ possesses a homogeneous in time statistics, in accordance with our general expectations. Note that $\chi \sim s/\gamma \gg 1$ as follows from Eq. (2.18). Keeping the main in χ contributions to the diagonal terms in Eq. (2.17) one obtains $\text{diag}(\partial_t \hat{\Delta} \hat{\Delta}^{-1}) = (-\chi\sigma, \chi\sigma, 0)$. Therefore in this approximation

$$\text{diag} \Delta = (e^{-\rho}, e^{\rho}, 1), \quad \partial_t \rho = \chi\sigma. \quad (2.19)$$

If $t \gg \lambda^{-1}$ then typically $\rho \sim \lambda t \gg 1$.

One concludes from the equations for $\zeta_1, \zeta_2, \zeta_3$ following from Eq. (2.17) that at $\lambda t \gg 1$ the variable ζ_1 is “frozen” at a level of order unity whereas the variables ζ_2, ζ_3 grow exponentially, they can be estimated as e^{ρ} . However, the combination $\zeta_1 \zeta_3 - \zeta_2$, entering $\hat{T}_R^{-1}$, is “frozen” at a level of order unity as well as ζ_1 . Based on the results obtained for the matrices $\hat{T}_L$, $\hat{\Delta}$, and $\hat{T}_R$ one finds eigen values of the matrix $\hat{W}$. In the main approximation in λ/s we obtain $W_1 \sim e^{\rho}$, $W_2 \sim 1$, $W_3 \sim e^{-\rho}$. The expressions together with Eq. (2.19) lead to the relation

$$\lambda \equiv \langle \partial_t \rho \rangle = \langle \chi \sigma \rangle, \quad (2.20)$$

where averaging is performed over the velocity statistics.

Since the variable ρ is an integral over time of a random quantity with homogeneous in time statistics, see Eqs. (2.18, 2.19), the ρ -statistics possesses some universal features at $\lambda t \gg 1$. Namely, at the condition the probability distribution function (PDF) of ρ can be written in a self-similar form [32]

$$P(\rho) \propto \exp[-tS(\rho/t)], \quad (2.21)$$

where S is the so-called Kramer function (or entropy function). The expression (2.21) is a manifestation of PDF's for the so-called intensive variables (see, e.g., [33]). The expression (2.21) implies that relative fluctuations of ρ diminish as t grows.

Let us consider moments of the divergence of close Lagrangian trajectories in our random flow. The equation governing a separation between the trajectories $\Delta \mathbf{R}$ is $\partial_t \Delta \mathbf{R}_j = \Sigma_{ji} \Delta \mathbf{R}_i$, it can be obtained from Eq. (2.9) by putting $\xi \rightarrow 0$. A solution of the equation is $\Delta \mathbf{R}(t) =$

$\hat{W}\Delta\mathbf{R}(0)$. Therefore at $t \gg \lambda^{-1}$ we arrive at the estimation $\Delta R(t) \sim \Delta R(0)e^\rho$. Then the moments of $\Delta\mathbf{R}$ can be calculated in the saddle-point approximation (justified by the inequality $\lambda t \gg 1$):

$$\langle |\Delta\mathbf{R}|^n \rangle = \int d\rho P(\rho) |\Delta\mathbf{R}|^n \propto \exp(\lambda_n t), \quad (2.22)$$

$$\lambda_n = -S(\psi_n) + n\psi_n, \quad \text{where } S'(\psi_n) = n. \quad (2.23)$$

Thus, the exponents λ_n are determined by statistical properties of the Lagrangian trajectories. Note that the Lyapunov exponent λ can be formally expressed via λ_n as $\lambda = d\lambda_n/dn|_{n=0}$.

General statistical properties of the separation $\Delta\mathbf{R}$ for the random flow with strong average shear were established in Ref. [21], in context of the single polymer dynamics in such flow. One expects a strong intermittency of $\Delta R(t)$, it should be revealed in faster than linear growth of λ_n at large n , since the linear law $\lambda_n \propto n$ is characteristic of Gaussian statistics of $\Delta R(t)$.

III. CORRELATION FUNCTIONS

To find a time dependence of the magnetic field moments one has to perform an additional average of the expression (2.14) over space that is equivalent to averaging over the ρ statistics and statistics of initial magnetic fluctuations. Thus, the $2n$ -th moment of the magnetic field induction is written as

$$\langle \mathbf{B}^{2n}(t) \rangle = \int d\rho P(\rho) [\mathbf{B}^{2n}(t)]. \quad (3.1)$$

In our approximation, $W_1 \sim e^\rho$, $W_2 \sim 1$ and, therefore, in the diffusionless regime $B(t) \sim e^\rho B_0$, whereas in the diffusion regime one obtains $|\mathbf{B}|^2 \sim \mathcal{B}_0^2(l/r_d)e^\rho$, as follows from Eq. (2.12). Substituting the expressions into Eq. (3.1) and integrating over ρ (in the saddle-point approximation) we find $\gamma_n = \lambda_{2n}$ for the diffusionless regime and $\gamma_n = \lambda_n$ for the diffusion regime. Thus, we related the dynamo increments (growth rates) introduced by Eq. (2.2) to the statistical properties of the flow. Our results can be summarized in terms of the estimations

$$\begin{aligned} \langle |\mathbf{B}(t)|^{2n} \rangle &\sim \exp(\lambda_{2n}t) \mathcal{B}_0^{2n}, \quad t < \lambda^{-1} \ln(l/r_d), \quad (3.2) \\ \langle |\mathbf{B}(t)|^{2n} \rangle &\sim (l/r_d)^n \exp(\lambda_n t) \mathcal{B}_0^{2n}, \quad t > \lambda^{-1} \ln(l/r_d). \end{aligned}$$

The main contribution to the moments $\langle \mathbf{B}^{2n}(t) \rangle$ is associated with the component B_x of the magnetic induction directed along the velocity of the shear flow, see Fig. 1. Let us turn to moments of the component B_y directed along the gradient of the shear flow, $\langle B_y^{2n} \rangle$. The moments are much smaller than the moments $\langle \mathbf{B}^{2n}(t) \rangle$, the smallness is caused by the strong shear flow. One finds from Eqs. (2.15, 2.19) that $[B_x^2(t)] = \chi^2 [B_y^2(t)]$. Thus, the variable χ is a measure of the magnetic field anisotropy, χ^{-1} determines the tilt angle ϕ of the magnetic blobs to the shear velocity, see Fig. 1. Since the

variable χ possesses homogeneous in time statistics, the factor χ^{-2} does not produce a difference in the growth rates, that is the both moments, $\langle B_x^{2n} \rangle$ and $\langle B_y^{2n} \rangle$, are proportional to the same exponent $\exp(\gamma_n t)$. However, the prefactors at the exponents are different. To find the difference in the prefactors is not enough to know statistical properties of the variable ρ that determine the exponent. Generally, one should know a mutual probability distribution of the variables $\rho(t)$ and $\chi(t)$ that is quite complicated object depending on details of the flow dynamics. However, one can establish an estimation for typical fluctuations $\chi \sim s/\lambda$ that follows from Eqs. (2.18, 2.20). Therefore, say, $[B_x^2(t)] \sim (s^2/\lambda^2) [B_y^2(t)]$.

There is a question concerning moments of the third component of the magnetic induction, $\langle B_z^{2n} \rangle$. To analyze their behavior one should take into account the components of the matrix $\hat{T}_L$, that we ignored at investigating B_x and B_y . Then we conclude that the time dependence of $\langle B_z^{2n} \rangle$ is characterized by the same exponents $\exp(\gamma_n t)$ both at the diffusionless and diffusion stages. As to prefactors, they depend on details of the flow statistics.

A. Pair Correlations

Let us consider the simultaneous magnetic field pair correlation function

$$F_{ij}(t, \mathbf{r}) = \langle B_i(t, \mathbf{r}_1 + \mathbf{r}) B_j(t, \mathbf{r}_1) \rangle. \quad (3.3)$$

Here angular brackets mean, as previously, averaging over space (that is an integration over $\mathbf{r}_1$ with the inverse volume as a factor). We assume statistical homogeneity in space of both the velocity and the initial magnetic field fluctuations, that is why the space average (3.3) characterizes the magnetic field correlations in the whole volume. We consider the case $r \ll \eta$, then we can use the smooth flow approximation.

Again, we start from the presentation (2.5). Then, analogously to the second moment, the pair correlation function (3.3) can be written as $F_{ij}(t) = \langle [W_{ik} \mathcal{B}_k[\mathbf{R}(0)] W'_{jl} \mathcal{B}_l[\mathbf{R}'(0)]] \rangle$, where the trajectories $\mathbf{R}$ and $\mathbf{R}'$ terminate at the points $\mathbf{r}_1 + \mathbf{r}$ and $\mathbf{r}_1$, respectively, at time t . Then one obtains

$$F_{ij}(t) = \langle [W_{ik}(t) W'_{jl}(t) \mathcal{F}_{kl}[\Delta\mathbf{R}(0)]] \rangle, \quad (3.4)$$

where $\mathcal{F}_{ij}$ is the initial (at $t = 0$) pair correlation function of the magnetic field fluctuations and $\Delta\mathbf{R} = \mathbf{R} - \mathbf{R}'$. The correlation length l of $\mathcal{F}$ is smaller than η , that is why we can consider $|\Delta\mathbf{R}| < \eta$. Then both evolution matrices in the expression (3.4) can be taken at the same point $\mathbf{R}$. Averaging in Eq. (3.4) can be treated as averaging over the velocity statistics.

The difference $\Delta\mathbf{R}$ satisfies the same equation (2.9) provided $|\Delta\mathbf{R}| \ll \eta$. However, now we are interested in the solution with the final condition $\Delta\mathbf{R} = \mathbf{r}$. Such

solution is written as

$$\Delta \mathbf{R}(t') = \hat{W}(t') \hat{W}^{-1}(t) \mathbf{r} - \hat{W}(t') \int_{t'}^t dt_1 \hat{W}^{-1}(t_1) [\boldsymbol{\xi}(t_1) - \boldsymbol{\xi}'(t_1)], \quad (3.5)$$

instead of Eq. (2.10). We immediately conclude from Eq. (3.5) that the pair correlation function coincides with the second moment if $r \lesssim r_d$. Therefore further we examine the case $r \gg r_d$ where the second term in Eq. (3.5) is negligible and we find

$$\Delta \mathbf{R}(0) = \hat{W}^{-1}(t) \mathbf{r}. \quad (3.6)$$

To be more precise, the expression (3.6) is correct provided $y \sim r \gg r_d$, that is implied below.

There are two different regimes for the pair correlation function. If $t < \lambda^{-1} \ln(l/r)$ then $\Delta \mathbf{R}(0)$ is typically less than l , the regime exists if $r \ll l$. In this case the two Lagrangian trajectories, $\mathbf{R}$ and $\mathbf{R}'$ remains typically within the correlation radius l at $t = 0$ and the behavior of the expression (3.4) is insensitive to the separation r . Therefore the pair correlation function F_{ij} practically coincides with the single-point average $\langle B_i B_j \rangle$ in this regime and, consequently, its time dependence is determined by the increment $\gamma = \lambda_2$.

If $t > \lambda^{-1} \ln(l/r)$ then $\Delta \mathbf{R}(0)$ is typically larger than l and only rare events where $|\Delta \mathbf{R}(0)| < l$ contribute to the correlation function. Using the representation $\hat{W} = \hat{T}_L \hat{\Delta} \hat{T}_R$ one obtains from Eq. (3.6) an expression $\Delta \mathbf{R}(0) \approx e^{\rho}(r_x - \chi r_y)$, where r_x and r_y are coordinates of the separation $\mathbf{r}$. A probability that the quantity is less or of the order of l is estimated as $e^{-\rho} l/r$ (provided $r_x \sim r_y \sim r$), that is an interval of values of χ where $\Delta \mathbf{R}(0) < l$. Therefore $B_i B_j \sim \mathcal{B}_0^2 e^{\rho} l/r$ and, consequently, $F(t) \sim \mathcal{B}_0^2 \exp(\lambda_1 t) l/r$.

Let us collect the obtained results

$$\begin{aligned} F(t) &\sim \mathcal{B}_0^2 \exp(\lambda_2 t), & t < \lambda^{-1} \ln(l/r) \\ F(t) &\sim \mathcal{B}_0^2 \exp(\lambda_1 t) l/r, & t > \lambda^{-1} \ln(l/r), \end{aligned} \quad (3.7)$$

where the inequality $r_d \ll r \ll l$ is assumed. Thus, the pair correlation function is governed by the same exponents as the second moment. In addition, we find an r -dependence of the pair correlation function. Note that the expression (3.7) turns into the expression (3.2) for the second moment at $r \sim r_d$ as it should be.

Returning to the expression (3.4) we conclude that a difference between the pair correlation function F_{ij} and the moments $\langle B_i B_j \rangle$ is solely in the behavior of $\Delta \mathbf{R}$. Therefore relations between components of F_{ij} controlled by the evolution matrices in Eq. (3.4) is the same as for the moments $\langle B_i B_j \rangle$, say, $F_{yy} \sim (\lambda/s)^2 F_{xx}$.

B. Mellin transform

It is instructive to examine the Mellin transform of the pair correlation function, the analysis reveals its scaling

properties. We define the Mellin transform as

$$\tilde{F}(t, k) = \int_0^\infty \frac{dr}{r} \left(\frac{r}{l} \right)^{-ik} F(t, r), \quad (3.8)$$

where a direction of the radius-vector $\mathbf{r}$ is implied to be fixed. Due to smoothness of the velocity field, different harmonics $\tilde{F}(t, k)$ evolve independently, being represented as a sum of exponents characterizing different structures of $\tilde{F}_{ij}$. At times $t \gg \lambda^{-1}$ only the principal exponent survives, that is $\tilde{F}(k, \varphi) \propto \exp[\gamma(k)t]$. In order to return back to the real space one should perform the inverse Mellin transform

$$F(t, r) = \int_{-\infty}^\infty \frac{dk}{2\pi} \exp \left[-ik \ln \frac{l}{r} + \gamma(k)t \right] \tilde{F}(k). \quad (3.9)$$

The quantity $\tilde{F}(k)$ is determined by the initial magnetic field fluctuations, correlated on the scale l . That is why we incorporated the quantity into the relations (3.8, 3.9).

Some words about analytical properties of $\tilde{F}(k)$. We assume that at $r > l$ the initial pair correlation function $\mathcal{F}(r)$ decreases fast as r grows. Then the integral (3.8) converges if $\text{Im } k > 0$. Therefore $\tilde{F}(k)$ is analytical in the upper k -semiplane. Besides, the integral diverges (at small r) as $k \rightarrow 0$. Therefore singularities of $\tilde{F}(k)$ lie in the lower k -semiplane, starting from the point $k = 0$. A character of the singularities depends on analytical properties of the initial function $\mathcal{F}(r)$. If it is analytical in r then one expects that $\tilde{F}(k)$ has a series of poles along the lower imaginary semiaxis, and the first one lies at $k = 0$. Note that in accordance with general rules the integration contour in Eq. (3.9) should go above the first singular point $k = 0$.

Now we can draw some general conclusions taking into account that $\gamma(k) \sim \lambda$. If $\ln(l/r) > \lambda t$ then the integral (3.9) is determined by a narrow vicinity of the point $k = 0$. Then $F(t, r) \propto \exp[\gamma(0)t]$ and we identify $\gamma(0)$ and λ_2 . If $\ln(l/r) < \lambda t$ then the integral (3.9) can be calculated in the saddle-point approximation. To find the saddle point one should shift the integration contour into the upper semiplane to reach the saddle-point $k = iq_*$, where q_* determines the minimal value of $\gamma(iq)$ at $q > 0$. Indeed, the increment $\gamma(iq)$ is real and therefore the point $k = iq_*$ is a solution of the equation $d\gamma/dk = 0$ giving an extremum of the exponent in Eq. (3.9). Then $F(t, r) \propto (l/r)^{q_*} \exp[\gamma(iq_*)t]$, and we identify $\gamma(iq_*)$ with λ_1 , see Eq. (3.7). Note that in accordance with the asymptotic law (3.7) q_* should be equal to unity, $q_* = 1$.

C. High Order Correlations

Here we consider higher order correlation functions of the magnetic field (2.7). One obtains expressions like Eq. (3.5) for separations $\Delta \mathbf{R}$ between the points

$\mathbf{R}_1(0), \dots, \mathbf{R}_{2n}(0)$ that are needed to calculate F_{2n} in accordance with Eq. (2.5). If $\lambda t < \ln(l/|\Delta\mathbf{r}|)$ for all separations between the points $\mathbf{r}_1, \dots, \mathbf{r}_{2n}$ then all separations $\Delta\mathbf{R}(0)$ are less than l . In this situation we arrive at the same expression $F_{2n} \sim \mathcal{B}_0^{2n} \langle e^{2n\rho} \rangle$ as for the $2n$ -th moment, and we conclude that $F_{2n} \propto \exp(\lambda_{2n}t)$, see Eq. (3.2).

Now we turn to the case where $\lambda t > \ln(l/r)$ where all separations $\Delta\mathbf{r}$ are assumed to be of the same order. Let us consider first the geometry where all the points $\mathbf{r}_1, \dots, \mathbf{r}_{2n}$ lie on a line, that is all vectors $\mathbf{r}_\alpha - \mathbf{r}_\beta$ have the same directions, and one can write $\Delta\mathbf{r} \sim \mathbf{r}$, where $\mathbf{r}$ is one of the separations. Then one arrives at the estimation $|\Delta\mathbf{R}| \sim e^\rho(r_x - \chi r_y)$, similar to the pair correlation function. Thus, one obtains the same probability $\sim e^{-\rho}l/r$ that the separations $|\Delta\mathbf{R}(0)| \lesssim l$. Then we find $F_{2n} \sim \mathcal{B}_0^{2n} \langle e^{(2n-1)\rho} \rangle l/r$, where the factor $e^{2n\rho}$ originates from the product of the matrices $\hat{W}$, appearing in accordance with the expression (2.5). Averaging the expression, one obtains $F_{2n}(t) \sim \mathcal{B}_0^{2n} \exp(\lambda_{2n-1}t)l/r$. Let us stress that the growth rates here are different from the growth rates of the corresponding moments.

However, the above expression is correct only if $t < \lambda^{-1} \ln(l/r_d)$. For larger t the diffusion contributions to the differences $\Delta_\alpha \mathbf{R}$ become relevant, see Eq. (3.5). Then, manipulating by χ , it is possible to make less than l only one difference among $\Delta\mathbf{R}(0)$. After that all other differences acquire typically values of the order of $r_d e^\rho$, and a probability that a difference is smaller than l is estimated as $(l/r_d)e^{-\rho}$. Thus we conclude $F_{2n} \sim \langle (l/r)(l/r_d)^{n-1} e^{n\rho} \rangle \propto \exp(\lambda_n t)$.

The same results (upto combinatoric factors) are true for a collinear geometry where the set $\mathbf{r}_1, \dots, \mathbf{r}_{2n}$ is separated into n pairs with parallel vectors $\mathbf{r}_\alpha - \mathbf{r}_\beta$ characterizing the pairs. Then the corresponding differences $\Delta\mathbf{R}$ behave like previously and the same arguments are correct. Let summarize our results for the collinear geometry

$$\begin{aligned} F_{2n}(t) &\sim \mathcal{B}_0^{2n} \exp(\lambda_{2n}t), \quad t < \lambda^{-1} \ln(l/r); \\ F_{2n}(t) &\sim \mathcal{B}_0^{2n} \exp(\lambda_{2n-1}t)l/r, \quad \lambda^{-1} \ln(l/r) < t < \lambda^{-1} \ln(l/r_d); \\ F_{2n}(t) &\sim \mathcal{B}_0^{2n} (l/r)(l/r_d)^{n-1} \exp(\lambda_n t), \quad t > \lambda^{-1} \ln(l/r_d). \end{aligned} \quad (3.10)$$

It is a generalization of the expressions (3.7) for the pair correlation function. Note also, that, as for the pair correlation function, in the collinear geometry the estimate $B_y \sim (\lambda/s)B_x$ determines relations between different components of F_{2n} whereas z -components need a separate investigation.

If the collinear geometry is destroyed, then it is impossible to put all the separations $|\Delta\mathbf{R}(0)|$ inside the scale l at any χ , if $t > \lambda^{-1} \ln(l/r)$. In this situation a behavior of the correlation function F_{2n} is non-universal being sensitive to details of the initial space distribution of $\mathcal{B}$. Any case, a value of F_{2n} in the non-collinear geometry is much less than in the collinear one. The situation resembles one realized for the randomly advected passive

scalar on scales larger than the pumping length [36]. We conclude that at $t > \lambda^{-1} \ln(l/r)$ correlations of the magnetic field are concentrated near collinear geometries, decaying away from the geometries. The decaying length is estimated as $le^{-\lambda t}$ at $t < \lambda^{-1} \ln(l/r_d)$ and as r_d at $t > \lambda^{-1} \ln(l/r_d)$.

IV. SHORT CORRELATED FLOW

Here we consider a strong steady shear flow complemented by a random component shortly correlated in time. The case admits an analytical solution and can be, therefore, used to verify our general assertions and predictions. In addition, the case is naturally realized since the strong shear destroys correlations of the random component, that is why we expect that the short-correlated case is widely spread in real flows.

In the short-correlated case the matrix of the velocity gradients $\hat{\sigma}$ describing the random component of the flow has to be treated as white noise, that is a variable δ -correlated in time. In the isotropic case one obtains the following tensorial structure

$$\langle \sigma_{ik}(t_1) \sigma_{jn}(t_2) \rangle = D(4\delta_{ij}\delta_{kn} - \delta_{ik}\delta_{jn} - \delta_{in}\delta_{jk})\delta(t_1 - t_2), \quad (4.1)$$

where the factor D characterizes strength of the random flow and the numerical factor is introduced as in Ref. [35]. However, as we already argued, the only relevant component of the random velocity gradients matrix in the case $\lambda \ll s$ (that is a manifestation of the random flow weakness) is $\sigma \equiv \sigma_{yx}$. We characterize its statistical properties by the expression

$$\langle \sigma(t_1) \sigma(t_2) \rangle = 4D\delta(t_1 - t_2), \quad (4.2)$$

formally coinciding with Eq. (4.1) for the yx -component. Other components of $\hat{\sigma}$ can have correlation functions different from the expression (4.1). The random component can be treated as weaker than the steady shear flow provided $D \ll s$.

Statistical properties of the separation between close Lagrangian trajectories $\Delta\mathbf{R}(t)$ in the random smooth flow with strong shear component in the short-correlated case are investigated in Ref. [37] (in the context of polymer dynamics). We present here results obtained in the paper without a derivation. Note that our variable χ is related to the tilt angle ϕ of Ref. [37] as $\chi = \cot \phi$, $\chi \approx \phi^{-1}$ in the case of small tilt angles, we are interested in. An expression for the Lyapunov exponent found in Ref. [37] is

$$\lambda = \frac{\sqrt{\pi} 3^{1/3}}{\Gamma(1/6)} D^{1/3} s^{2/3}. \quad (4.3)$$

Thus, the condition $s \gg D$ guarantees the inequality $\lambda \ll s$, indeed. Note also that $\lambda \gg D$ and that $\lambda \rightarrow 0$ as $D \rightarrow 0$. The last property is a natural consequence of zero value of the Lyapunov exponent for a purely shear flow.

For the short-correlated case, it is possible to find the exponents λ_n characterizing growth rates of the moments of $\Delta \mathbf{R}(t)$, see Eq. (2.22), if $n \gg 1$. Then a saddle-point (instanton) approximation in the functional space can be used, see Ref. [34], that leads to

$$\gamma_n = \frac{3}{2^{5/3}} n^{4/3} D^{1/3} s^{2/3} \sim \lambda n^{4/3}, \quad (4.4)$$

see Ref. [37]. The non-linear dependence of λ_n on n , $\lambda_n \propto n^{4/3}$, signals about strong intermittency of the flow. Note, that in our anisotropic case the growth rates λ_n increase as n grows slower than in the isotropic case, where for the short-correlated flow $\lambda_n \propto n^2$, see Ref. [15].

A. Pair Correlation Function

Next, we examine the two-point simultaneous correlation function (3.3) for the short-correlated in time velocity field. In the case it is possible to derive a closed equation for the correlation function (see, e.g., Ref. [12]). We study here the function at scales that are much larger than the diffusion scale r_d (but much smaller than the velocity correlation length η). Then it is possible to neglect diffusion effects and we omit all terms with the noise ξ in subsequent relations.

Let us shortly explain a derivation of the equation. First of all, it follows from the definition (2.6) that

$$\hat{W}(t) = T \exp(\hat{\Xi}) \hat{W}(t - \tau), \quad \hat{\Xi} = \int_{t-\tau}^t dt' \hat{\Sigma}, \quad (4.5)$$

where τ is an arbitrary time (less than t). We choose τ to be much smaller than λ^{-1} but much larger than the velocity correlation time (it is possible for the short-correlated flow). Then $\hat{\Xi}$ is a small factor, though the two factors in Eq. (4.5) can be treated as statistically independent. Substituting the expression (4.5) into Eq. (3.4), expanding the result into a series over $\hat{\Xi}$ (upto the second order) and averaging the result inside the time interval $(t - \tau, t)$ in accordance with Eq. (4.2), one obtains a variation of F_{ij} at passing from $t - \tau$ to t . Since the variation is small, it can be rewritten in terms of a differential equation.

Assuming the isotropic correlation function of the fluctuations (4.1) one obtains the following equation

$$\begin{aligned} \partial_t F_{ij} = & -2sy\partial_x F_{ij} + sF_{iy}\delta_{jx} + sF_{yj}\delta_{ix} + \\ & + 4D \left[\delta_{ij} F_{kk} - \frac{1}{2} F_{ij} - r_k \partial_j F_{ik} - r_k \partial_i F_{kj} + \right. \\ & \left. + \frac{1}{2} (r \nabla) F_{ij} + \frac{1}{2} r^2 \nabla^2 F_{ij} - \frac{1}{4} r_m r_n \partial_m \partial_n F_{ij} \right]. \end{aligned} \quad (4.6)$$

In the absence of the shear (at $s = 0$) the system of equations (4.6) leads to a closed equation for the trace of the correlation function $H = F_{kk}$:

$$\partial_t H = D (10H + 6r\partial_r H + r^2\partial_r^2 H).$$

The equation coincides with one presented in Ref. [14] (for the case of the scaling exponent $\xi = 2$ and zero forcing).

Now we rule out irrelevant terms in Eq. (4.6) using the following properties: s is much larger than D , characteristic value of x is much larger than that of y and, respectively, $\partial_y \gg \partial_x$. Then we can keep in Eq. (4.6) solely the terms originating from σ_{yx} . The resulting equation leads to a closed system of equations for the three components, F_{xx} , F_{xy} , F_{yy} , of the pair correlation function:

$$\begin{aligned} \partial_t F_{xx} = & -2sy\partial_x F_{xx} + 2sF_{xy} + 4Dx^2\partial_y^2 F_{xx}, \\ \partial_t F_{xy} = & -2sy\partial_x F_{xy} + sF_{yy} + 2Dx^2\partial_y^2 F_{xy} - 4Dx\partial_y F_{xx}, \\ \partial_t F_{yy} = & -2sy\partial_x F_{yy} + 2Dx^2\partial_y^2 F_{yy} - 8Dx\partial_y F_{xy} + DF_{xx}. \end{aligned} \quad (4.7)$$

Further, we use the dimensionless time $T = (8Ds^2)^{1/3}t$ and introduce the designations $f = F_{xx}$, $g = (s/D)^{1/3}F_{xy}$ and $h = (s/D)^{2/3}F_{yy}$.

Let us investigate a special case of coinciding points. At $\mathbf{r} = 0$ all terms with derivatives drop from the equations (4.7) and they take the form

$$\partial_T \begin{pmatrix} f \\ g \\ h \end{pmatrix} = \begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & 1/2 \\ 2 & 0 & 0 \end{pmatrix} \begin{pmatrix} f \\ g \\ h \end{pmatrix}. \quad (4.8)$$

A growing solution of the solution is

$$\begin{pmatrix} f \\ g \\ h \end{pmatrix}_{r=0} \propto \begin{pmatrix} 1 \\ 1 \\ 2 \end{pmatrix} e^T. \quad (4.9)$$

The behavior corresponds to the increment $\gamma = (8Ds^2)^{1/3}$ of the magnetic field second moment. From the other hand, the case corresponds to small r that is to the condition $\ln(l/r) > \lambda t$. Therefore $\gamma = \lambda_2$, and we conclude that $\lambda_2 = (8Ds^2)^{1/3}$ in our case.

It is convenient to pass into the ‘‘polar coordinates’’ ϱ, φ in the shear plane: $x = \varrho \cos \varphi$, $y = (D/s)^{1/3} \varrho \sin \varphi$. Then we perform the Mellin transform $f, g, h \rightarrow \tilde{f}, \tilde{g}, \tilde{h}$ in terms of ϱ and derive the equations for $\tilde{f}, \tilde{g}, \tilde{h}$ from the system (4.7). In terms of the quantity $q = -ik$ the equations are written as

$$\begin{aligned} \partial_T \tilde{f} = & \tilde{f}'' \cos^4 \varphi + \tilde{f}' [\sin^2 \varphi - 2(q+1) \sin \varphi \cos^3 \varphi] \\ & + \tilde{f} [q \sin \varphi \cos \varphi - q \cos^4 \varphi + (q^2 + q) \sin^2 \varphi \cos^2 \varphi] + \tilde{g}, \\ \partial_T \tilde{g} = & \tilde{g}'' \cos^4 \varphi + \tilde{g}' [\sin^2 \varphi - 2(q+1) \sin \varphi \cos^3 \varphi] + \\ & + \tilde{g} [q \sin \varphi \cos \varphi - q \cos^4 \varphi + (q^2 + q) \sin^2 \varphi \cos^2 \varphi] \\ & - 2 \cos^2 \varphi \tilde{f}' + 2q \sin \varphi \cos \varphi \tilde{f} + \frac{1}{2} \tilde{h}, \\ \partial_T \tilde{h} = & \tilde{h}'' \cos^4 \varphi + \tilde{h}' [\sin^2 \varphi - 2(q+1) \sin \varphi \cos^3 \varphi] + \\ & + \tilde{h} [q \sin \varphi \cos \varphi - q \cos^4 \varphi + (q^2 + q) \sin^2 \varphi \cos^2 \varphi] \\ & - 4 \cos^2 \varphi \tilde{g}' + 4q \sin \varphi \cos \varphi \tilde{g} + 2\tilde{f}, \end{aligned} \quad (4.10)$$

where prime designates a derivative over the angle φ .




FIG. 2: Increment of the Mellin transform of the pair correlation function on the imaginary axis.

Next, we study the time evolution of the system (4.10) numerically for different real values of q (so that $k = iq$ is imaginary) using implicit difference scheme on the interval $(-\pi/2, \pi/2)$ for φ with the periodic boundary conditions. We have chosen as initial conditions for f, g, h the same Gaussian functions centered near $\varphi = 0$ and with width of order unity. Then we extract the principal increment $\gamma(iq) = (8Ds^2)^{1/3}c(q)$ dominating the behavior of the system at $T \gg 1$. The dimensionless quantity c was extracted as

$$c = \frac{1}{T} \ln \frac{f(T_0 + T)}{f(T_0)},$$

where $T_0 + T$ is chosen to be large enough (near 30) and T_0 is introduced to exclude an influence of an initial transient process, we have chosen $T_0 = T$.

The quantity c is plotted as a function of q in Fig. 2. It turned out to be positive everywhere, having a minimum at $q = 1$, $c(1) = 0.435$. The value $q = 1$ is in accordance with Eq. (3.7) and the general analysis given in Section III. As we argued there, the minimum value of $\gamma(iq)$ determines λ_1 , that is $\lambda_1 = c(q_*)(8Ds^2)^{1/3}$. The value of c at $q = 0$ is $c = 1$, in accordance with Eq. (4.9) and general arguments given in Section III. Thus, the obtained results confirm our general assertions.

V. DISCUSSION

We have analyzed the kinematic dynamo stage when the small-scale fluctuations of the magnetic field grow in a steady shear flow complemented by relatively weak random velocity fluctuations. The weakness is characterized by the inequality $s \gg \lambda$ where s is the shear rate and λ is the Lyapunov exponent of the flow. Universal features we are established are revealed at times $t \gg \lambda^{-1}$. The shear makes the flow strongly anisotropic, which, paradoxically, simplifies the analysis of the dynamo phenomenon since

a single component of the random velocity gradient appears to be relevant. We analyzed the situation where the correlation length l of initial magnetic field fluctuations is less than the velocity correlation length η (that is the Kolmogorov length for developed turbulence). Probably, the smallness of l is not crucial for our scheme since small scales of the magnetic field distribution in space are inevitably produced by the hydrodynamic motion.

Let us underline that in the main approximation in λ/s our problem is reduced to a purely two-dimensional velocity field (with components along the shear velocity and along its gradient). We have proved an existence of the dynamo in this case (that is the exponential growth of the magnetic field moments). The result obviously contradicts the statement of Refs. [38–40] (Zeldovich theorem) that there cannot be the magnetic dynamo in two-dimensional flows. We assert that this statement is wrong and the error is in ignoring the third component B_3 of the magnetic induction (perpendicular to the velocity plane). The third component satisfies the passive scalar equation and, consequently, decays exponentially. However, B_3 cannot be ignored in the divergentless condition $\nabla \mathbf{B} = 0$ since the characteristic scale of the magnetic field along direction of its growth increases faster than the magnetic field itself. One can check that all the terms in $\nabla \mathbf{B} = \partial_x B_x + \partial_y B_y + \partial_z B_z$ decay with the same exponent, and therefore the condition $\nabla \mathbf{B} = 0$ leads to an effectively divergent in-plane magnetic field. The dynamo effect is not forbidden for such field. A detailed analysis of the discrepancy will be published elsewhere. An existence of the dynamo effect for two-dimensional flows is a subject of numerical verification.

For our general conclusions, we do not specify statistical properties of the random flow exploring only its smoothness at scales less than the velocity correlation length η . Then it is possible to relate the kinematic growth rates γ_n of the magnetic field, see Eq. (2.2), to intrinsic characteristics of the flow characterizing divergence of close Lagrangian trajectories, see Eq. (2.22). We find that $\gamma_n = \lambda_{2n}$ in the diffusionless regime and $\gamma_n = \lambda_n$ in the diffusion regime. We also related the anisotropy degree of the magnetic field to the same intrinsic characteristics of the flow. Thus, the main features of the magnetic field statistics (including its intermittency) are dictated by the flow statistics. Note that our general scheme can be applied without essential modifications to the statistically isotropic flows or to random flows with other types of anisotropy.

We established main features of the magnetic field correlation functions. The pair correlation function behaves like the second moment at small separations r , and increase with the growth rate, characteristic of the diffusion regime, at larger r , it is proportional to $1/r$ there. As to higher order correlation functions, the situation is more complicated. As small time t they behave as corresponding moments. However, at larger time, $t > \lambda^{-1} \ln(l/r)$, correlations are peaked near the collinear geometry (where $2n$ points are separated into n pairs

with parallel separations) and there is an intermediate asymptotic regime when the correlation functions grow with rates that do not coincide with the growth rates of the moments. Then, at times $t > \lambda^{-1} \ln(l/r_d)$ the correlation function grows with the same exponent, as the corresponding moment in the diffusion regime. The scaling behavior of the correlation functions in the collinear regime is $\propto 1/r$. Correlations decay fast at deviation from the collinear geometry. That reflects a complicated spacial structure of the magnetic field that is strongly correlated for special geometries produced by affine geometrical transformations from initial magnetic fluctuations.

Our general assertions can be verified by solving the model with short correlated in time fluctuating component, that admits a series of analytical results. The non-linear n -dependence of the growth rates γ_n , $\gamma_n \propto n^{4/3}$, at large n signals about strong intermittency of the magnetic field. Therefore only rare events concentrated in a restricted fraction of space contribute to high moments

of the magnetic field. We analyzed in detail the pair correlation function of the magnetic field for the model, the analysis confirms all our general assertions, including scaling behavior in different regimes.

The ideology and the analytical approach developed in our work can be expanded to the dynamics of polymer solutions possessing elasticity that is described similarly to the magnetic field. Moving along these lines, we hope to clarify some aspects of the so-called elastic turbulence [22–24] that are still not explained.

Acknowledgments

We thank A. Chernykh, M. Chertkov, G. Falkovich, and S. S. Vergeles for helpful discussions and remarks. The work was partially supported by RFBR under grant no. 09-02-01346-a and by MES of RF under FTP “Kadry”.

-
- [1] Monin A. S. and Yaglom A. M., Statistical Fluid Mechanics, Vol. 1 and 2 MIT Press Cambridge Mass, 1975.
 - [2] Frisch U., Turbulence: the Legacy of A. N. Kolmogorov, Cambridge University Press New York, 1995.
 - [3] Vainshtein S. and Zeldovich Ya., Origin of magnetic fields in astrophysics, Sov. Phys. Usp. **15** 159 (1972).
 - [4] Moffatt H. K., Magnetic Field Generation in Electrically Conducting Fluids, Cambridge University Press, 1978.
 - [5] Parker E. N., Cosmic magnetic fields their origin and activity, Clarendon Press Oxford, 1979.
 - [6] Zeldovich Ya. B., Ruzmaikin A. A., and Sokolov D. D., The Almighty Chance, World Scientific, 1990.
 - [7] Childress S. and Gilbert A., Stretch Twist Fold: The Fast Dynamo, Springer-Verlag Berlin, 1995.
 - [8] Rüdiger G. and Hollerbach R., The Magnetic Universe: Geophysical and Astrophysical Dynamo Theory, Wiley, 2004.
 - [9] Fluid Dynamics and Dynamos in Astrophysics and Geophysics, eds. Soward A. M., Jones Ch. A., Hughes D. W., and Weiss N. O. CRC, 2005.
 - [10] Vishniac E. T., Lazarian A., and Cho J., Problems and Progress in Astrophysical Dynamos Springer, 2008.
 - [11] Schekochihin A. A., Cowley S. C., Taylor S. F., Maron L., and McWilliams J. C., Simulations of the small-scale turbulent dynamo, Astrophysical Journal **612** 276-307 (2004).
 - [12] Falkovich G., Gawedzki K., and Vergassola M., Particles and fields in fluid turbulence, Rev. Mod. Phys. **73** 913 (2001).
 - [13] Kraichnan R. and Nagarajan S., Growth of turbulent magnetic fields. Phys. Fluids **10** 859 (1967).
 - [14] Kazantsev A. P., Enhancement of a magnetic field by a conducting fluid, Zh. Eksp. Teor. Fiz. **53**, 18061813 (1967) [Sov. Phys. JETP **26** 1031 (1968)].
 - [15] Chertkov M., Falkovich G., Kolokolov I., and Vergassola M., Small-scale Turbulent Dynamo, Phys. Rev. Lett. **83** 4065-4068 (1999).
 - [16] Urpin V., Kinematic turbulent dynamo in a shear flow, J. Geophysical and Astrophysical Fluid Dynamics **95** 269-284 (2001).
 - [17] Bird R. B., Curtiss C. F., Armstrong R. C., and Hassager O., Dynamics of Polymeric Liquids, 2nd ed., Wiley, New York, 1987.
 - [18] Doi M. and Edwards S. F., The Theory of Polymer Dynamics Oxford University Press, 1988.
 - [19] Lumley J. L., On the solution of equations describing small scale deformation, Symp. Math. **9** 315 (1972).
 - [20] Balkovsky E., Fouxon A., and Lebedev V., Turbulent Dynamics of Polymer Solutions, Phys. Rev. Lett. **84** 4765-4768 (2000).
 - [21] Chertkov M., Kolokolov I., Lebedev V., and Turitsyn K., Polymer statistics in a random flow with mean shear, J. Fluid Mech. **531** 251-260 (2005).
 - [22] Groisman A. and Steinberg V., Elastic Turbulence in a Polymer Solution Flow, Nature **405** 53 (2000).
 - [23] Groisman A. and Steinberg V., Stretching of Polymers in a Random Three-dimensional Flow, Phys. Rev. Lett. **86** 934 (2001).
 - [24] Groisman A. and Steinberg V., Efficient Mixing at Low Reynolds Numbers Using Polymer Additives, Nature **410** 905 (2001).
 - [25] Kogan V. R., Kolokolov I. V., and Lebedev V. V., Kinematic magnetic dynamo in a random flow with strong average shear, J. Phys. A: Math. Theor. **43** 182001 (2010).
 - [26] Landau L. D. and Lifshitz E. M., Course of theoretical physics, Electrodynamics of Continuous Media, Pergamon, 1984.
 - [27] Chertkov M., Kolokolov I., and Vergassola M., Inverse cascade and intermittency of passive scalar in one-dimensional smooth flow, Phys. Rev. E **56** 5483-5499 (1997).
 - [28] Pope S. B., Lagrangian pdf Methods for Turbulent Flows, Ann. Rev. Fluid Mech. **26** 23 (1994).
 - [29] Furstenberg H., Noncommuting Random Products, Trans. Am. Math. Soc. **108** 377 (1963).
 - [30] Goldhirsch I., Sulem P.-L., and Orszag S. A., Stability

- and Lyapunov Stability of Dynamical Systems, *Physica D* **27** 311 (1987).
- [31] Balkovsky E. and Fouxon A., Universal Long-time Properties of Lagrangian Statistics in the Batchelor Regime and Their Application to the Passive Scalar Problem, *Phys. Rev. E* **60** 4164 (1999).
 - [32] Ellis R., *Entropy Large Deviations and Statistical Mechanics*, Springer Verlag, 1985.
 - [33] Landau L. D. and Lifshitz E. M., *Course of theoretical physics, Statistical Physics part 1*, Pergamon Press, 1980.
 - [34] Falkovich G., Kolokolov I., Lebedev V., and Migdal A., Instantons and Intermittency, *Phys. Rev. E* **54** 4896 (1996).
 - [35] Chertkov M., Falkovich G., Kolokolov V., Lebedev V., Normal and anomalous scaling of the fourth-order correlation function of a randomly advected passive scalar, *Phys. Rev. E* **52** 4924-4941 (1995).
 - [36] Balkovsky E., Falkovich G., Lebedev V., and Lysiansky M., Large-scale properties of passive scalar advection, *Phys. Fluids* **11** 2269-2279 (1999).
 - [37] Turitsyn K. S., Polymer dynamics in chaotic flows with a strong shear component, *ZhETF* **132** 746 (2007) [*JETP* **105** 655 (2007)].
 - [38] Zeldovich Ya. B., The magnetic field in the two-dimensional motion of a conducting turbulent fluid, *ZhETF* **31** 154 (1956) [*Sov. Phys. JETP* **4** 460 (1957)].
 - [39] Zeldovich Ya. B. and Ruzmaikin A. A., Magnetic field of a conducting fluid in two-dimensional motion, *ZhETF* **78** 980 (1980) [*Sov. Phys. JETP* **51** 493 (1980)].
 - [40] Zeldovich Ya. B., Ruzmaikin A. A., Molchanov S. A., and Sokolov D. D., Kinematic Dynamo Problem in a Linear Velocity Field, *J. Fluid Mech.* **144** 1 (1984).